



UNIVERSIDAD
DE SANTIAGO
DE CHILE

BOLETÍN LATINOAMERICANO Y DEL CARIBE
DE PLANTAS MEDICINALES Y AROMÁTICAS
17 (1): 30 - 35 (2018)
© / ISSN 0717 7917 / www.blacpma.usach.cl



Artículo / Article

Effects of essential oils from two *Lippia* species on growth of phytopathogenic fungi

[Efecto de los aceites esenciales de dos especies de *Lippia* sobre el crecimiento de hongos fitopatógenos]

Laura E. Leal^{1,2}, Alejandro A. Alarcón¹, Pablo Ortega-Baes², Fany Cayo¹ & Rosana Alarcón¹

¹Laboratorio de Productos Naturales. Facultad de Ciencias Naturales, Universidad Nacional de Salta (UNSa), 4400 Salta, Argentina.

²LABIBO (CONICET). Facultad de Ciencias Naturales, Universidad Nacional de Salta (UNSa), 4400 Salta, Argentina.

Contactos / Contacts: Rosana ALARCÓN - E-mail address: ralarcon@unsa.edu.ar

Abstract: Chemical characterization of the essential oils of two *Lippia* species by GC-MS and NMR spectroscopy revealed that limonene (84.3%) and β -caryophyllene (6.1%) were the most abundant components in *Lippia turbinata* while (6S,7S,10S)-trans-davanone (99.1%) predominated in *Lippia integrifolia*. Antifungal activity of the essential oils was determined by headspace volatile exposure assay against the fungal phytopathogenic *Sclerotinia sclerotiorum*, *Sclerotium rolfsii* and *Rhizoctonia solani*. The essential oil of *L. turbinata* showed potent antifungal activity against the panel of fungi tested while that the oil of *L. integrifolia* significantly inhibited the mycelial growth of *S. rolfsii* and *R. solani*.

Keywords: *Lippia turbinata*, *Lippia integrifolia*, essential oils, chemical composition, antifungal activity

Resumen: La caracterización química de los aceites esenciales de dos especies de *Lippia* por cromatografía gaseosa-espectrometría de masas (CG-EM) y espectroscopia de RMN reveló que limoneno (84,3%) y β -cariofileno (6,1%) fueron los componentes más abundantes de *Lippia turbinata* mientras que (6S,7S,10S)-trans-davanona (99,1%) predominó en *Lippia integrifolia*. La actividad antifúngica de los aceites esenciales se determinó por el ensayo de exposición a los vapores frente a los hongos fitopatógenos *Sclerotinia sclerotiorum*, *Sclerotium rolfsii* y *Rhizoctonia solani*. El aceite esencial de *L. turbinata* mostró una potente actividad antifúngica frente al panel de hongos ensayados, mientras que el aceite de *L. integrifolia* inhibió significativamente el crecimiento micelial de *S. rolfsii* y *R. solani*.

Palabras clave: *Lippia turbinata*, *Lippia integrifolia*, aceites esenciales, composición química, actividad antifúngica

Recibido | Received: February 20, 2017

Aceptado | Accepted: May 4, 2017

Aceptado en versión corregida | Accepted in revised form: June 5, 2017

Publicado en línea | Published online: January 30, 2018

Declaración de intereses | Declaration of interests: Financial support for this work by the Consejo de Investigación de la Universidad Nacional de Salta is gratefully acknowledged. LEL holds a research fellowship from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

Este artículo puede ser citado como / This article must be cited as: LE Leal, AA Alarcón, P Ortega-Baes, Fany Cayo, R Alarcón. 2017. Effects of essential oils from two *Lippia* species on growth of phytopathogenic fungi. *Bol Latinoam Caribe Plant Med Aromat* 17 (1): 30 – 35.

INTRODUCTION

Lippia is a genus which belongs to the Verbenaceae family; it is represented by more than 200 species, distributed mainly in South and Central America, and Africa (Sanders, 2001). Several species of the genus *Lippia* are used in folk medicine mostly in dermatological, gastrointestinal and respiratory affections. In addition, most of these species are used as food condiment and flavoring (Pascual *et al.*, 2001). In Argentina, the two most widely used native species are *Lippia integrifolia* (Griseb.) Hieron ("incayuyo", "té del inca" or "inca yerba"), and *Lippia turbinata* Griseb. ("Poleo", "té del país" or "té criollo"). In recent years, these species have become subject of scientific interest in view of other potential uses of its essential oils, for the most part, as antimicrobial (Bluma & Etcheverry, 2008; González & Marioli, 2010; Lima *et al.*, 2011; Passone *et al.*, 2013; Pérez-Zamora *et al.*, 2016), and antioxidant agents (Quiroga *et al.*, 2013; Barbieri *et al.*, 2016).

To our knowledge, the biological activities of the oils of *L. turbinata* and *L. integrifolia* collected in Salta province, Argentina, have not yet been explored. Therefore, the present work was undertaken with the main objective to investigate the effects of the essential oils on the radial growth of three phytopathogenic fungi, *Sclerotinia sclerotiorum*, *Sclerotium rolfsii* and *Rhizoctonia solani*, by headspace volatile exposure assay. Previous reports on *L. integrifolia* and *L. turbinata* EOs have indicated antifungal activity (Bluma & Etcheverry, 2008; Lima *et al.*, 2011; Passone *et al.*, 2013), but they were performed on different fungal species from those assayed herein.

MATERIAL AND METHODS

Plant Material

Aerial parts of *L. integrifolia* and *L. turbinata* were collected at the flowering stage in Salta, Argentina. Voucher specimens were deposited at the Museo de la Facultad de Ciencias Naturales, Universidad Nacional de Salta, Salta, Argentina. Plant identification was made according to voucher specimen numbers MCNS 11880 for *L. integrifolia* and MCNS 11879 for *L. turbinata*.

Isolation of the essential oils

Dried aerial parts were subjected to hydrodistillation for 4 h in a Clevenger type apparatus. The essential oils obtained were stored at 4° C until further analysis.

GC-FID and GC/MS analysis

GC/MS analyses of the EOs were performed with a Perkin-Elmer Series Clarus 600 gas chromatograph equipped with a flame ionization detector (FID) and Clarus 600 T mass spectrometer detector, and a DB-5 MS fused-silica cap. column (60 m, 0.25 mm, 0.25 µm). The oven temperature was programmed isothermal at 60° C for 5 min, then rising from 60 a 240° C at 4° C/min, and isothermal at 240° C for 10 min; injector and detector temp. 250° C; transferline temp., 200° C; carrier gas, He (49.6 psi); column head pressure, 15 psi. The split-injection mode was selected, and the ionization was carried out in the mass spectrometer under vacuum by electron impact (ionization energy, 70 eV). The chromatograms were acquired (scan mode, m/z 25-350) with a scan time of 0.2 s and an inter-scan time of 0.1 s. Diluted samples (1/100, v/v, in EtOH) of 1.0 µL were injected in the split mode (split ratio 50/1).

The constituents were identified by comparing their retention indices and mass spectra with those found on the NIST 02 library and in the literature (Adams, 2001). Retention indices were determined in relation to a homologous series of n-alkanes (C₇ – C₁₆ on DB-5) under the same operating conditions. The relative amounts of individual components were calculated on the GC-peak area (FID response) without using correction factors.

NMR Spectroscopy

NMR spectra were recorded on a Bruker Avance 400 (¹H at 400 MHz and ¹³C at 100 MHz) spectrometer with TMS as internal reference. The sample was dissolved in CDCl₃. ¹H NMR and ¹³C NMR analyses were performed on the whole sample, without any previous fractionation. The identification was based on comparison of the signals in the oil spectrum with reference spectra compiled in the laboratory spectral library and the literature (Kubeczka & Formacek, 2002; Skakovskii *et al.*, 2010; Marcial *et al.*, 2016).

Limonene: ¹H NMR (400 MHz, CDCl₃): δ 5.40 (m, H-6), 4.70 (br s, H-10), 1.97 (m, H-2), 1.81-1.76 (m, H-3), 2.10 (m, H-4), 2.05-1.93 (m, H-5), 1.65 (s, H-7), 1.73 (br s, H-9). ¹³C NMR (100 MHz, CDCl₃): δ 133.7 (s, C-1), 30.6 (t, C-2), 27.9 (t, C-3), 41.1 (d, C-4), 30.8 (t, C-5), 120.6 (d, C-6), 23.2 (q, C-7), 150.3 (s, C-8), 20.9 (q, C-9), 108.3 (t, C-10).

(6S,7S,10S)-trans-davanone: ¹H NMR (400 MHz, CDCl₃): δ 1.75 (d, *J*= 1.1, H-1), 5.33 (m, H-3), 3.21

(dd, $J = 6.9, 18.1$, H-4a), 3.28 (dd, $J = 6.9, 18.1$, H-4b), 2.70 (dq, $J = 6.9, 8.5$, H-6), 4.05 (ddd, $J = 6.9, 8.4$, H-7), 1.66 (m, H-8a), 1.70 (m, H-8b), 1.71 (m, H-9a), 1.86 (m, H-9b), 5.80 (dd, $J = 10.6, 17.2$, H-11), 4.95 (dd, $J = 9.1, 1.6$, H-12a), 5.13 (dd, $J = 17.3, 1.6$, H-12b), 1.62 (br s, H-13), 1.01 (d, $J = 6.9$, H-14), 1.29 (s, H-15). ^{13}C NMR (100 MHz, CDCl_3): δ 25.7 (q, C-1), 135.0 (s, C-2), 116.1 (d, C-3), 42.7 (t, C-4), 211.9 (s, C-5), 51.2 (d, C-6), 80.5 (d, C-7), 29.3 (t, C-8), 36.8 (t, C-9), 83.0 (s, C-10), 143.6 (d, C-11), 111.3 (t, C-12), 18.1 (q, C-13), 12.8 (q, C-14), 27.2 (q, C-15).

Antifungal Activity Assay

The fungi tests were field isolates maintained on potato dextrose agar (PDA) plates (*Sclerotinia sclerotiorum*, *Sclerotium rolfsii*, *Rhizoctonia solani*). Inoculated plates were sealed with parafilm and incubated at 20° C until confluent growth across the agar surface occurred. Plates were then stored at 4° C. Antifungal activity was determined by headspace volatile exposure assay.

To agar plates containing 5 mL PDA in glass Petri dishes ($\varnothing = 50$ mm) placed with the lid upside down was added essential oil in a small jar plugged with cotton having no direct contact with the PDA agar plates (Vilela *et al.*, 2009). Oil doses were 0 (control), 10 and 20 μL /Petri dish. At the center of each dish a 5 mm diameter disk of fungal species cultivated for not less than 7 days was placed. The plates were sealed with polyethylene film and incubated in the dark at $22 \pm 2^\circ \text{C}$. Five replicates were prepared for each treatment. The oil samples and controls were randomly arranged. The colony diameter was measured after 3 days for *S. sclerotiorum* and *S. rolfsii*, and after 7 days for *R.*

solani. The percentage inhibition of growth was calculated for each colony according to the following formula: % Inhibition = $[(C-T)/C] \times 100$, where C is the average diameter of the colonies of control, in mm, and T is the average diameter of the colonies developed in each treatment. The fungicidal or fungistatic activity of the vapor of *L. turbinata* originating from different application concentrations was confirmed by transferring half of the mycelial plug onto fresh PDA agar after 3 days for *S. sclerotiorum* and *S. rolfsii*, and after 7 days for *R. solani*. The activity of EO was regarded as fungicidal if 100% growth inhibition after initial exposure was maintained following transfer of the inoculated plug to fresh agar.

Statistical analysis

The statistical significance of assay results was determined by variance analysis with the Infostat Programme. Differences between means were tested by Kruskal-Wallis test; differences between experiment and control were significant with a value of $p \leq 0.05$.

RESULTS AND DISCUSSION

Hydrodistillation of the dried aerial parts of *L. integrifolia* and *L. turbinata* gave yellow essential oils in yields of 0.76% and 0.32% (w/w) respectively. Table 1 shows the components of EOs. The major component of *L. integrifolia* was davanone (99.1%), while the essential oil of *L. turbinata* was characterized by a high percentage of monoterpenes, limonene (84.3%) being the main component.




Figure 1
Chromatogram of *Lippia integrifolia* essential oil

A recent study reported that essential oil composition of *L. integrifolia* collected from different locations in Argentina had different chemical constituents. According to this report, five chemotypes were observed: *trans*-davanone chemotype, lippifolienone chemotype, β -davanone-2-ol chemotype, spathulenol/bicyclogermacrene chemotype, and *trans*-nerolidol chemotype (Marcial *et al.*, 2016). In this study, the high percentage of davanone (99.1%) clearly indicates that the plant belongs to *trans*-davanone chemotype. In addition, the specific optical rotation recorded for essential oil of *L. integrifolia* { $[\alpha]_D$ (20° C) + 65.9 (c 1.0 in

CHCl₃) } was in agreement with those reported for (6*S*,7*S*,10*S*)-*trans*-davanone (Marcial *et al.*, 2016).

In this work, at the doses tested (Table 2), the essential oil of *L. integrifolia* showed significant antifungal activities (inhibition % > 60%) against *R. solani* and *S. rolfsii*, while not affecting the growth of *S. sclerotiorum*.

In a previous study, essential oil of *L. integrifolia* with borneol, terpinen-4-ol and lippifolienone as major constituents has shown, *in vitro*, an interesting spectrum of antifungal properties (Lima *et al.*, 2011). However we did not find in the literature any report of antifungal properties of essential oils of *trans*-davanone chemotype.

Table 1

Composition of the essential oils extracted from aerial parts of *Lippia turbinata* and *Lippia integrifolia*.

Components	RI ^a	Peak area (%)		Identification
		<i>L. integrifolia</i>	<i>L. turbinata</i>	
Davanone	1588	99.1	-	MS, RI, ¹ H and ¹³ C NMR
Camphene	954	-	Tr	MS, RI
Limonene	1034	-	84.3	MS, RI, ¹ H and ¹³ C NMR
Camphor	1146	-	1.5	MS, RI
Carvone	1245	-	1.1	MS, RI
Piperitenone oxide	1369	-	1.3	MS, RI
β -caryophyllene	1419	-	6.1	MS, RI
Total		99.1	94.3	

^a Retention indices on a DB-5 column; tr = trace amount (< 0.05%)




Figure 2
Chromatogram of *Lippia turbinata* essential oil

Table 2
Growth inhibition of fungal species by essential oils of *Lippia turbinata* and *Lippia integrifolia*.

Essential Oils	Dose (μL of oil/petri dish)	<i>Sclerotium rolfii</i>	<i>Sclerotinia sclerotiorum</i>	<i>Rhizoctonia solani</i>
		Growth mm	Growth mm	Growth mm
<i>L. turbinata</i>	0	45.0 ± 0.0	45.0 ± 0.0	45.0 ± 0.0
	10	0.0 ± 0.0* (100)	0.0 ± 0.0 a* (100)	0.0 ± 0.0 (100)
	20	0.0 ± 0.0* (100)	0.0 ± 0.0* (100)	0.0 ± 0.0 (100)
<i>L. integrifolia</i>	0	45.0 ± 0.0	45.0 ± 0.0	44.8 ± 0.4
	10	15.6 ± 8.7* (65.3)	38.4 ± 7.2 (14.7)	10.4 ± 1.8* (76.9)
	20	12.7 ± 7.2* (71.8)	33.1 ± 17.8 (26.4)	9.6 ± 3.5* (78.7)

The growth of fungal species is given as mean ± SD of five replicates. *Significant differences, compared to the control for $p \leq 0.05$, according to Kruskal-Wallis's test. Figures in parenthesis indicate percent inhibition

Moreover, the essential oil of *L. turbinata* caused complete growth inhibition of all the pathogens tested via a fungicidal mode. It is highly probable that the antifungal activity of *L. turbinata* essential oil is attributed to its high limonene content (84.3%). This compound evidences fungicidal activity against *Trichophyton rubrum*, inducing changes in cell membrane integrity and metabolic activity (Chee et al., 2009). Recently, it has been reported that limonene may show antifungal activity by inhibition of pectin methyl esterase (PME) and cellulase enzymes (Marei et al., 2012). In previous studies, other authors (Bluma & Etcheverry, 2008; Passone et al., 2012; Passone et al., 2013) have reported antifungal properties for this oil, finding piperitenone oxide (48.6%) and limonene (24.5%) as major components, while in our study, piperitenone oxide only represented 1.6% of the oil. They evaluated the antifungal activity against *Aspergillus flavus* and *Aspergillus parasiticus*. In general, EO composition showed significant chemodiversity according to extraction method and geographical origin (Bakkali et al., 2008), showing the importance of analyzing plants from different locations.

CONCLUSIONS

The composition of our *L. turbinata* sample agrees with that in previous reports of plants collected from Cordoba province, Argentina (Quiroga et al., 2013), regarding the presence of limonene as a major constituent (% > 75). On the other hand, our findings

are in agreement with those of the previous chemical composition analysis of plants of *L. integrifolia* from the same province (Marcial et al., 2016). It should be noted that in our oil, content of (6*S*,7*S*,10*S*)-trans-davanone (99.1%) is the highest reported so far for this ketone in any other essential oil.

In our study, *L. turbinata* essential oil was highly effective in vapour phase against *S. sclerotiorum*, *S. rolfii* and *R. solani*, and its activity was higher than *L. integrifolia* EO. In addition, this is the first report on the biological activities of trans-davanone chemotype from *L. integrifolia*.

ACKNOWLEDGEMENTS

Financial support for this work by the Consejo de Investigación de la Universidad Nacional de Salta is gratefully acknowledged. LEL holds a research fellowship from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

REFERENCES

- Adams P. 2001. **Identification of Essential Oils components by Gas Chromatography/Quadrupole Mass Spectroscopy**. Allured Publishing Corporation, Illinois, USA.
- Bakkali F, Averbeck S, Averbeck D, Idaomar M. 2008. Biological effects of essential oils- A review. **Food Chem Toxicol** 46: 446 - 475.
- Barbieri N, Costamagna M, Gilabert M, Perotti M, Schuff C, Isla MI, Benavente A. 2016.

- Antioxidant activity and chemical composition of essential oils of three aromatic plants from La Rioja province. **Pharm Biol** 54: 168 - 173.
- Bluma R, Etcheverry M, 2008. Application of essential oil in maize grain: impact on *Aspergillus* section *Flavi* growth parameters and aflatoxin accumulation. **Food Microbiol** 25: 324 - 334.
- Chee HY, Kim H, Lee MH. 2009. In vitro Antifungal Activity of Limonene against *Trichophyton rubrum*. **Mycobiology** 37: 243 - 246.
- González MJ, Marioli JM. 2010. Antibacterial activity of water extracts and essential oils of various aromatic plants against *Paenibacillus larvae*, the causative agent of American Foulbrood. **J Invertebr Pathol** 104: 209 - 213.
- Kubeczka KH, Formacek V. 2002. **Essential Oils Analysis by Capillary Gas Chromatography and Carbon-13 NMR Spectroscopy**. Wiley, New York, USA.
- Lima B, López S, Luna L, Agüero MB, Aragón L, Tapia A, Zacchino S, López ML, Zygodlo J, Feresin GE. 2011. Essential oils of medicinal plants from the central andes of Argentina: Chemical composition, and antifungal, antibacterial, and insect-repellent activities. **Chem Biodivers** 8: 924 - 936.
- Marcial G, De Lampasona MP, Vega MI, Lizarraga E, Viturro CI, Slanis A, Juárez MA, Elechosa MA, Catalán CAN. 2016. Intraspecific variation in essential oil composition of the medicinal plant *Lippia integrifolia* (Verbenaceae). Evidence for five chemotypes. **Phytochemistry** 122: 203 - 212.
- Marei GIK, Abdel Rasoul MA, Abdelgaleil SAM. 2012. Comparative antifungal activities and biochemical effects of monoterpenes on plant pathogenic fungi. **Pestic Biochem Physiol** 103: 56 - 61.
- Pascual ME, Slowing K, Carretero E, Sánchez Mata D, Villar A. 2001. *Lippia*: Traditional uses, chemistry and pharmacology: A review. **J Ethnopharmacol** 76: 201 - 214.
- Passone MA, Girardi NS, Ferrand CA, Etcheverry M. 2012. In vitro evaluation of five essential oils as botanical fungitoxicans for the protection of stored peanut of *Aspergillus flavus* and *A. parasiticus* contamination. **Int Biodeterior Biodegrad** 70: 82 - 88.
- Passone MA, Girardi NS, Etcheverry M. 2013. Antifungal and antiaflatoxigenic activity by vapor contact of three essential oils, and effects of environmental factors on their efficacy. **LWT - Food Sci Technol** 53: 434 - 444.
- Pérez-Zamora MC, Torres CA, Aguado MI, Bela AJ, Nuñez MB, Bregni C. 2016. Antibacterial activity of essential oils of *Aloysia polystachya* and *Lippia turbinata* (Verbenaceae). **Bol Latinoam Caribe Plant Med Aromat** 15: 199 - 205.
- Quiroga PR, Grosso NR, Lante A, Lomolino G, Zygodlo JA, Nepote V. 2013. Chemical composition, antioxidant activity and antilipase activity of *Origanum vulgare* and *Lippia turbinata* essential oils. **Int J Food Sci Technol** 48: 642 - 649.
- Sanders RW. 2001. The genera of Verbenaceae in the southeastern United States. **Harv Pap Bot** 5: 303 - 358.
- Skakovskii ED, Kiselev WP, Tychinskaya LY, Schutova AG, Gonsharova LW, Spiridowish EW, Bovdey NA, Kiselev PA, Gaidukevich OA. 2010. Characterization of the essential oil of *Agastache rugosa* by NMR spectroscopy. **J Appl Spectrosc** 77: 329 - 334.
- Vilela GR, de Almeida GS, D'Arce MABR, Moraes MHD, Brito JO, da Silva MFGF, Silva SC, de Stefano Piedade SM, Calori-Domingues MA, da Gloria EM. 2009. Activity of essential oil and its major compound, 1,8-cineole, from *Eucalyptus globulus* Labill., against the storage fungi *Aspergillus flavus* Link and *Aspergillus parasiticus* Speare. **J Stored Prod Res** 45: 108 - 111.